

Anti-Probable mitochondrial-processing peptidase subunit alpha-1, mitochondrial antibody

Catalog: PHY1410A

Product Information

Description:	Rabbit polyclonal antibody
Background:	Mitochondrial ubiquinol-cytochrome c oxidoreductase is also called complex III
	and it accepts electron from quinol and transferred to cytochrome c. In
	Arabidopsis mitochondria, ubiquinol-cytochrome c oxidoreductase contains 10
	subunits. MPP alpha (AT1G51980) is one of the subunit of Complex III.
Synonyms:	AT1G51980
Immunogen:	KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from
	Arabidopsis thaliana AT1G51980.
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70°C as supplied.
	6 months, -20 to -70°C under sterile conditions after reconstitution.
	1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the
	temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	54 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 80-99% homologues with the sequence in
	Brassica napus, Brassica rapa, Glycine max, Vitis vinifera, Medicago


Research Use Only



truncatula, Cucumis sativus.

The sequence of the synthetic peptide used for immunization is 93% homologues with the sequence in Alpha-MPP 2 (AT3G16480). For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 15 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 40 kDa. Electrophoresis: 12% SDS-PAGE Transfer: blotting to NC (nitrocellulose) membrane for 1 h. Blocking: 5% skim milk at RT or 4°C for 1 h. Primary antibody: 1:1000 dilution overnight at 4°C. Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only