

## Anti-Protein ELC antibody

Catalog: PHY3393A

## **Product Information**

Description:	Rabbit polyclonal antibody
Background:	ELC is the component of the ESCRT-I complex (endosomal sorting complex
	required for transport I), a regulator of vesicular trafficking process. It is
	required for the sorting of endocytic ubiquitinated cargos into multivesicular
	bodies (MVBs). And it may control nuclear division through the microtubule
	cytoskeleton.
Synonyms:	ELC, VPS23A, ATELC, VACUOLAR PROTEIN SORTING 23A
Immunogen:	KLH-conjugated synthetic peptide (15 aa from Central section) derived from
	Arabidopsis thaliana ELC (AT3G12400).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
<b>Reconstitution:</b>	Reconstitution with 150 μl of 0.01 M sterile PBS.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70°C as supplied.
	6 months, -20 to -70°C under sterile conditions after reconstitution.
	1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the
	temperature recommended above.

## **Application Information**


<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected/apparent MW:	45 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used for
	immunization is 100% homologues with the sequence in Brassica
	napus and 80-99% homologues with the sequence in Brassica rapa



Vitis vinifera, Gossypium raimondii, Nicotiana tabacum, Cucumis sativus, Glycine max, Medicago truncatula. For more species homologues information, please contact tech

support at tech@phytoab.com.

## **Application Example**



Recom: 2.5 ng, 15 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 44 kDa. Electrophoresis: 12% SDS-PAGE Transfer: blotting to NC (nitrocellulose) membrane for 1 h. Blocking: 5% skim milk at RT or 4°C for 1 h. Primary antibody: 1:1000 dilution overnight at 4°C. Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only

PhytoAB Inc.