

## Anti-Alternative NAD(P)H-ubiquinone oxidoreductase C1, chloroplastic/mitochondrial antibody

Catalog: PHY3413A

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** NDC1 catalyzes the penultimate step in phylloquinone (vitamin K1)

biosynthesis.

**Synonyms:** NDC1, NAD(P)H DEHYDROGENASE C1

**Immunogen:** KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana NDC1 (AT5G08740).

Form: Lyophilized

**Quantity:** 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to  $-70^{\circ}$ C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected / apparent MW:** 57 kDa

**Predicted Reactivity:** Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in Solanum

tuberosum, Glycine max, Solanum lycopersicum, Vitis vinifera,

Gossypium raimondii, Medicago truncatula, Nicotiana tabacum, Vitis vinifera, , and 80-99% homologues with the sequence in Brassica


Research Use Only



napus, Cucumis sativus, Populus trichocarpa, Spinacia oleracea, Sorghum bicolor, Brassica rapa.

For more species homologues information, please contact tech support at <a href="mailto:tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 2.5 ng, 15 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 47 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.