

Anti-BTB/POZ domain and ankyrin repeat-containing protein NPR2 antibody

Catalog: PHY4469S

Product Information

Description:	Rabbit polyclonal antibody
Background:	Similar to Ankyrin repeat BTB/POZ domain-containing protein. (Os01t0767900-01)
Synonyms:	NPR2, Arabidopsis NPR1 homolog 2, NPR1 homologue 2, NPR1-like 2, non-expressor of pathogenesis-related gene2, nonexpresser of PR genes 2/OsNH2, OsNPR2, OsNPR2/NH2
Immunogen:	KLH-conjugated synthetic peptide (17 aa from C terminal section) derived from <i>Orvza sativa</i> NPR2 (Os01g0767900).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum
Reconstitution:	Reconstitution with 50µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	68 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in <i>Zea mays</i> , <i>Sorghum bicolor</i> , and 80-99% homologues with the sequence

Research Use Only

in *Triticum aestivum*, *Hordeum vulgare*, *Setaria viridis*, *Panicum virgatum*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.