

Anti-Transcription factor TCP4, C-terminal antibody

Catalog: PHY3595A

Product Information

Description:	Rabbit polyclonal antibody	
Background:	TCP4 is a transcription factor. It is regulated by miR319 and involved in	
	heterchr	onic regulation of leaf differentiation.
Synonyms:	TCP4, N	IATERNAL EFFECT EMBRYO ARREST 35, MEE35, TCP FAMILY
	TRANSC	CRIPTION FACTOR 4
Immunogen:	KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from	
	Arabidop	osis thaliana TCP4 (AT3G15030).
Form:	Lyophilized	
Quantity:	150 μg	
Purification:	Immunogen affinity purified	
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS.	
	"Note: pl	lease spin tube briefly prior to opening it to avoid any losses that might
	occur fro	om lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ \!\! \mathbb C$ as supplied.	
	6 months, -20 to -70 $^\circ\!\!\!\mathrm{C}$ under sterile conditions after reconstitution.	
	1 month	, 2 to 8 $^\circ \!\!\!\! \mathbb{C}$ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\!\mathbb{C}$. Upon receipt, store it immediately at the	
	temperature recommended above.	
Application Info	ormatio	n
Recommended Dilution:		Western Plat (1:1000 1:2000)
Recommended Dilu	uon.	Western Blot (1:1000-1:2000)
		Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent	• RA\A/•	46 kDa
Expected / apparent MW:		
Predicted Reactivity	/:	Among species analyzed, the sequence of the synthetic peptide used
		for immunization is 80-99% homologues with the sequence in
		Brassica rapa, Brassica napus.
		For more species homologues information, please contact tech
		support at <u>tech@phytoab.com</u> .

Research Use Only