

Anti-Protein SUPPRESSOR OF K(+) TRANSPORT GROWTH DEFECT 1 antibody

Catalog: PHY0864A

Product Information

Description: Rabbit polyclonal antibody

Background: SKD1 (Suppressor of K+ Transport Growth Defect 1) is a homolog

of Vps4p/SKD1 which localized to the cytoplasm and to multivesicular

endosomes. It is involved in the transport of biosynthetic membrane proteins

from the prevacuolar/endosomal compartment to the vacuole.

Synonyms: SKD1, ATSKD1, SUPPRESSOR OF K+ TRANSPORT GROWTH DEFECT1,

VACUOLAR PROTEIN SORTING 4, VPS4

Immunogen: KLH-conjugated synthetic peptide (19 aa from N terminal section) derived from

Arabidopsis thaliana SKD1 (AT2G27600).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 49 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 80-99% homologues with the sequence in Vitis



vinifera, Glycine max, Cucumis sativus, Oryza sativa.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with

CCD camera.