

Anti-SPINDLY antibody

Product Information

Description:	Rabbit polyclonal antibody	
Background:	SPY is a N-acetyl glucosamine transferase that may glycosylate other	
	molecules involved in GA signaling. SPY acts as both a repressor of GA	
	responses and as a positive regulation of cytokinin signalling. It may be	
	involved in reducing ROS accumulation in response to stress.	
Synonyms:	SPY, SPINDLY	
Immunogen:	Recombinant protein (1-149) of SPY derived from Arabidopsis	
	thaliana AT3G11540.	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Serum	
Reconstitution:	Reconstitution with 150 μl of ste <mark>r</mark> ile water.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ \! \mathbb C$ as supplied.	
	6 months, -20 to -70 $^\circ\!\!\!\!\!^\circ$ under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^{\circ}$ C under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4 $^\circ\!{ m C}$. Upon receipt, store it immediately at the	
	temperature recommended above.	

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)	
	Note: Optimal dilutions/concentrations should be determined by the	
	end user.	
Expected/apparent MW:	101 kDa	
Predicted Reactivity:	For more species homologues information, please contact tech	
	support at tech@phytoab.com.	

Research Use Only



Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 18 kDa.
Electrophoresis: 12% SDS-PAGE
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.



Research Use Only