

Anti-Protein TIC110, chloroplastic antibody

Catalog: PHY1364A

Product Information

Description:	Rabbit polyclonal antibody
Background:	TIC110 is a protein thought to be a part of the translocon at the chloroplast
	inner envelope. It is involved in protein import into the chloroplast and
	chloroplast biogenesis.
Synonyms:	TIC110, ARABIDOPSIS THALIANA TRANSLOCON AT THE INNER
	ENVELOPE MEMBRANE OF CHLOROPLASTS 110, ATTIC110,
	TRANSLOCON AT THE INNER ENVELOPE MEMBRANE OF
	CHLOROPLASTS 110.
Immunogen:	KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from
	Arabidopsis thaliana TIC110 (AT1G06950).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 μl of 0.01 M sterile PBS.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ \!\! \mathbb{C}$ as supplied.
	6 months, -20 to -70 $^\circ$ C under sterile conditions after reconstitution.
	1 month, 2 to 8 $^\circ$ C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the
	temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	112 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 100% homologues with the sequence in <i>Brassica</i>

Research Use Only



napus, Brassica rapa, and 80-99% homologues with the sequence in Nicotiana tabacum, Glycine max, Medicago truncatula, Cucumis sativus, Gossypium raimondii. For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 54 kDa.
Electrophoresis: 12% SDS-PAGE
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).
Detection: using chemiluminescence substrate and image were captured with

CCD camera.

Research Use Only