

Anti-EPIDERMAL PATTERNING FACTOR-like protein 9, C-terminal antibody

Catalog: PHY0990A

Product Information

Description: Rabbit polyclonal antibody

Background: ATEPFL9 is a cysteine-rich peptide, a secretory factor that is produced in the

mesophyll cells and acts on the epidermis to increase stomatal formation. Its mature form is a 45-aa peptide with three intramolecular disulfide bonds. It is proposed that ATEPFL9 increases stomatal number by competing with two

negative regulators of stomatal density, EPF1 and EPF2.

Synonyms: EPFL9, ATEPFL9, EPIDERMAL PATTERNING FACTOR LIKE-9, STOMAGEN

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana EPFL9 (AT4G12970).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4° C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 12 kDa




Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in Gossypium raimondii, Vitis vinifera, Cucumis sativus, Brassica rapa, Brassica napus, Glycine max, Populus trichocarpa, Medicago truncatula, Nicotiana tabacum, Solanum lycopersicum, Solanum tuberosum.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 58 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with

CCD camera.