

Anti-Metal transporter Nramp3 antibody

Catalog: PHY0709S

Product Information

Description:	Rabbit polyclonal antibody	
Background:	NRAMP3 is a member of the Nramp2 metal transporter family; like its homolog	
	Atnramp4 (AT5G67330), localized in vacuolar membrane.	
Synonyms:	NRAMP3, ATNRAMP3, NATURAL RESISTANCE-ASSOCIATED	
	MACROPHAGE PROTEIN 3	
Immunogen:	KLH-conjugated synthetic peptide (14 aa from Central section) derived from	
	Arabidopsis thaliana NRAMP3 (AT2G23150).	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Serum	
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .	
Reconstitution:	Reconstitution with 150µl of sterile water.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ\!{ m C}$ as supplied.	
	6 months, -20 to -70 $^\circ\!\!\!\!\!^\circ$ under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^{\circ}$ C under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4 $^\circ\!\!\!\mathrm{C}$. Upon receipt, store it immediately at the	
	temperature recommended above.	
Reference:	Viviane Lanquar, et al., Plant Physiology, April 2010, Vol. 152, pp. 1986–1999.	

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	56 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 80-99% homologues with the sequence in
	Brassica napus, Brassica rapa.

Research Use Only



For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 52 kDa.
Electrophoresis: 12% SDS-PAGE
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).
Detection: using chemiluminescence substrate and image were captured

with CCD camera.

Research Use Only