

Anti-PsbP domain-containing protein 1, chloroplastic antibody

Catalog: PHY3618A

Product Information

Description: Rabbit polyclonal antibody

Background: PPD1 is a nuclear-encoded thylakoid lumenal protein that is associated with

photosystem I (PSI) but is not an integral subunit of PSI. It is a PSI assembly factor that assists the proper folding and integration of PsaB and PsaA into the

thylakoid membrane. It is involved in the assembly of PSI reaction center

heterodimer core.

Synonyms: PPD1, PSBP-DOMAIN PROTEIN1

Immunogen: KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from

Arabidopsis thaliana PPD1 (AT4G15510).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 32 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 80-99% homologues with the sequence in



Brassica napus, Brassica rapa.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 40 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.