

Anti-Autophagy-related protein 8A/B/C, C-terminal antibody

Catalog: PHY4410S

Product Information

Description:	Rabbit polyclonal antibody
Background:	ATG8A/B/C are autophagy associated protein.
Synonyms:	ATG8A/B/C, OsATG8a/b/c
Immunogen:	KLH-conjugated synthetic peptide (16 aa from C terminal section but not the cleavage region of ATG4) derived from <i>Oryza sativa</i> ATG8A (Os07g0512200), ATG8B (Os04g0624000) and ATG8C (Os08g0191600).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum
	Peptide affinity form antibody available upon request at info@phytoab.com .
Reconstitution:	Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	14 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologous with the sequence in <i>Zea mays</i> , <i>Gossypium raimondii</i> , <i>Solanum tuberosum</i> , <i>Brassica rapa</i> , <i>Brassica napus</i> , <i>Hordeum vulgare</i> , <i>Vitis vinifera</i> , <i>Glycine max</i> ,

Research Use Only

Arabidopsis thaliana, Nicotiana tabacum, Solanum lycopersicum, Populus trichocarpa, Setaria viridis, Medicago truncatula, Panicum virgatum, Spinacia oleracea, Cucumis sativus, Sorghum bicolor, Triticum aestivum.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 46 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.