

Anti-Cre05.g242400, C-terminal antibody

Catalog: PHY5593S

Product Information

Description:	Rabbit polyclonal antibody
Background:	Cre05.g242400
Synonyms:	Cre05.g242400
Immunogen:	KLH-conjugated synthetic peptide (20 aa from C terminal section) derived from <i>Chlamydomonas reinhardtii</i> Cre05.g242400 .
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum Peptide affinity form antibody available upon request at info@phytoab.com .
Reconstitution:	Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	15 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in <i>Sorghum bicolor</i> , <i>Zea mays</i> , <i>Panicum virgatum</i> , <i>Physcomitrium patens</i> , <i>Gossypium raimondii</i> , <i>Solanum tuberosum</i> , <i>Solanum lycopersicum</i> , <i>Setaria viridis</i> , <i>Cucumis sativus</i> , <i>Spinacia oleracea</i> , <i>Oryza sativa</i> ,

Research Use Only

Triticum aestivum, Hordeum vulgare, Nicotiana tabacum, Vitis vinifera, Populus trichocarpa, Brassica rapa, Brassica napus, Arabidopsis thaliana, Glycine max, Medicago truncatula.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 53 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.