

## Anti-Protein ACTIVITY OF BC1 COMPLEX KINASE 1, chloroplastic antibody

Catalog: PHY3463A

## **Product Information**

Description:	Rabbit polyclonal antibody
Background:	ACDO1 belongs to protein kinase superfamily protein, it is required for
	photooxidative stress responses, including the induction of oxidative stress
	response genes to prevent photosystem II core and chlorophyll degradations.
Synonyms:	ACDO1, ABC1-LIKE KINASE 1, ABC1-LIKE KINASE RELATED TO
	CHLOROPHYLL DEGRADATION AND OXIDATIVE STRESS 1, ABC1K1,
	ATACDO1, PGR6, PROTON GRADIENT REGULATION 6
Immunogen:	KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from
	Arabidopsis thaliana ACDO1 (AT4G31390).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
<b>Reconstitution:</b>	Reconstitution with 150 µl of 0.01 M sterile PBS.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ C$ as supplied.
	6 months, -20 to -70 $^\circ$ C under sterile conditions after reconstitution.
	1 month, 2 to 8 $^\circ\!\mathrm{C}$ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}$ . Upon receipt, store it immediately at the
	temperature recommended above.

## **Application Information**

<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	77 kDa

PhytoAB Inc.




**Predicted Reactivity:** 

Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in *Brassica rapa*, *Brassica napus*.

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 42 kDa.
Electrophoresis: 12% SDS-PAGE
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).
Detection: using chemiluminescence substrate and image were

Research Use Only

PhytoAB Inc.