

Anti-GLYMA.06G110800 antibody

Catalog: PHY6524S

Product Information

Description:	Rabbit polyclonal antibody	
Background:	GLYMA.06G110800.	
Synonyms:	GLYMA.06G110800, GmNSP2	
Immunogen:	KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from	
	Glycine max GLYMA.06G110800.	
Form:	Lyophilized	
Quantity:	150 μg	
Purification:	Serum	
	Peptide affinity form antibody available upon request at info@phytoab.com.	
Reconstitution:	Reconstitution with 150 μl of sterile water.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ\!\mathrm{C}$ as supplied.	
	6 months, -20 to -70 $^\circ C$ under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^\circ \!$	
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the	
	temperature recommended above.	

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	54 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 80-99% homologues with the sequence in <i>Zea</i>
	mays, Hordeum vulgare, Oryza sativa, Setaria viridis, Triticum
	aestivum, Panicum virgatum, Sorghum bicolor, Vitis vinifera, Medicago
	truncatula.

PhytoAB Inc.



The sequence of the synthetic peptide used for immunization is 93% (14/15) homologues with the sequence in GLYMA.04G251900. For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



PHY6524S

Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 48 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4 °C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only