

## **Anti-Protein CYCLOPS antibody**

Catalog: PHY6514S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** GLYMA.01G149500 is a DNA-binding transcriptional activator.

Synonyms: GLYMA.01G149500, GmCYCLOPS-1

Immunogen: KLH-conjugated synthetic peptide (18 aa from C terminal section) derived from

Glycine max GLYMA.01G149500.

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 58 kDa

**Predicted Reactivity:** The sequence of the synthetic peptide used for immunization is 94%


(17/18) homologues with the sequence in GLYMA.09G213700. For more species homologues information, please contact tech

Tof more species nomologues information, please contact tec

support at tech@phytoab.com.



## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 47 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.