

Anti-BZIP domain-containing protein antibody

Catalog: PHY6512S

Product Information

Description:	Rabbit polyclonal antibody	
Background:	GmSTF3/4 are soybean TGACG-motif binding factors.	
Synonyms:	GmSTF3/4	
Immunogen:	KLH-conjugated synthetic peptide (14 aa from Central section) derived from	
	Glycine max GmSTF3 (Glyma.16G092700) and GmSTF4 (Glyma.03G081700).	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Serum	
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .	
Reconstitution:	Reconstitution with 150 µl of sterile water.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ C$ as supplied.	
	6 months, -20 to -70 $^{\circ}$ C under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^\circ C$ under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the	
	temperature recommended above.	

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	22 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 100% homologues with the sequence in <i>Medicago</i>
	<i>truncatula</i> , and 80-99% homologues with the sequence in <i>Zea may</i> s,
	Triticum aestivum, Oryza sativa, Sorghum bicolor, Hordeum vulgare,
	Spinacia oleracea, Setaria viridis, Panicum virgatum, Populus


PhytoAB Inc.



trichocarpa.

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 40 kDa. **Electrophoresis:** 12% SDS-PAGE **Transfer:** blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4°C for 1 h. **Primary antibody:** 1:1000 dilution overnight at 4°C. **Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000). **Detection:** using chemiluminescence substrate and image were

captured with CCD camera.

Research Use Only