

Anti-Chlorophyll a-b binding protein, chloroplastic antibody

Catalog: PHY5663S

Product Information

Description:	Rabbit polyclonal antibody
Background:	LHCB5
Synonyms:	LHCB5, Ihcb5
Immunogen:	KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from
	Chlamydomonas reinhardtii LHCB5 (Cre16.g673650).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .
Reconstitution:	Reconstitution with 150 µl of sterile water.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ \!\! \mathbb{C}$ as supplied.
	6 months, -20 to -70 $^\circ\!\!\!\mathrm{C}$ under sterile conditions after reconstitution.
	1 month, 2 to 8 $^\circ C$ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4° C. Upon receipt, store it immediately at the
	temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	31 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 80-99% homologues with the sequence in <i>Populus</i>
	trichocarpa, Glycine max, Gossypium raimondii, Vitis vinifera,
	Nicotiana tabacum, Solanum tuberosum, Solanum lycopersicum,

PhytoAB Inc.


Research Use Only



Cucumis sativus, Arabidopsis thaliana, Spinacia oleracea, Hordeum vulgare, Triticum aestivum, Setaria viridis, Zea mays, Sorghum bicolor, Panicum virgatum, Brassica rapa, Brassica napus, Medicago truncatula.

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



PHY5663S

Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa. **Electrophoresis:** 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4° C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only