

Anti-tRNA (cytosine(38)-C(5))-methyltransferase 2 antibody

Catalog: PHY3499A

Product Information

| Description: | Rabbit polyclonal antibody |
|------------------------|--|
| Background: | DNMT2 is a DNA methyltransferase homolog. Human Dnmt2 methylates |
| | tRNA-Asp and can methylate Arabidopsis tRNA-Asp in vitro. |
| Synonyms: | DNMT2, ATDNMT2, DNA METHYLTRANSFERASE 2, TRDMT1, DNA |
| | METHYLTRANSFERASE-2 |
| Immunogen: | KLH-conjugated synthetic peptide (14 aa from Central section) derived from |
| | Arabidopsis thaliana DNMT2 (AT5G25480). |
| Form: | Lyophilized |
| Quantity: | 150 µg |
| Purification: | Immunogen affinity purified |
| Reconstitution: | Reconstitution with 150 µl of 0.01 M sterile PBS. |
| | "Note: please spin tube briefly prior to opening it to avoid any losses that might |
| | occur from lyophilized material adhering to the cap or sides of the tube". |
| Stability & | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. |
| Storage: | 12 months from date of receipt, -20 to -70 $^\circ \! \mathbb{C}$ as supplied. |
| | 6 months, -20 to -70 $^\circ\!\!\!\!^\circ$ under sterile conditions after reconstitution. |
| | 1 month, 2 to 8 $^{\circ}$ C under sterile conditions after reconstitution. |
| Shipping: | The product is shipped at 4 $^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the |
| | temperature recommended above. |


Application Information

| Recommended Dilution: | Western Blot (1:1000-1:2000) |
|------------------------------|--|
| | Note: Optimal dilutions/concentrations should be determined by the |
| | end user. |
| Expected / apparent MW: | 44 kDa |
| Predicted Reactivity: | For more species homologues information, please contact tech |
| | support at tech@phytoab.com. |

Research Use Only



Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 47 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.



Research Use Only