

## **Anti-Transcription factor MYB75 antibody**

Catalog: PHY1193S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** MYB75 is a putative MYB domain containing transcription factor involved in

anthocyanin metabolism and radical scavenging. It is essential for the sucrose-mediated expression of the dihydroflavonol reductase gene. It

interacts with JAZ proteins to regulate anthocyanin accumulation.

Synonyms: MYB75, PAP1, ARABIDOPSIS THALIANA PRODUCTION OF ANTHOCYANIN

PIGMENT 1, ATMYB75, ATPAP1, MYB DOMAIN PROTEIN 75,

MYELOBLASTOSIS PROTEIN 75, PRODUCTION OF ANTHOCYANIN

PIGMENT 1, SIAA1, SUC-INDUCED ANTHOCYANIN ACCUMULATION 1

**Immunogen:** KLH-conjugated synthetic peptide (14 aa from C terminal section) derived from

Arabidopsis thaliana MYB75 (AT1G56650).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

**Reconstitution:** Reconstitution with 150µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability & Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4 °C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected/apparent MW: 28 kDa


Research Use Only



**Predicted Reactivity:** 

For more species homologues information, please contact tech support at <a href="tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.