

Anti-Probable zinc metalloprotease EGY1, chloroplastic antibody

Catalog: PHY2868S

Product Information


Description:	Rabbit polyclonal antibody
Background:	Membrane-associated and ATP-independent metalloprotease; EGY1 protein contains eight trans-membrane domains at its C-terminus, and carries out beta-casein degradation in an ATP-independent manner. EGY1 is required for development of both thylakoid grana and a well-organized lamellae system in chloroplast. Additionally, EGY1 is required for the accumulation of chlorophyll and chlorophyll a/b binding (CAB) proteins (both PS I and PS II) in chloroplast membranes, and for grana formation and normal chloroplast development. Loss of EGY1 function also has an effect on endodermal plastid biogenesis. Mutant studies suggest that EGY1 is involved in the regulation of nuclear gene expression response to ammonium stress and interacts with ABA signaling.
Synonyms:	EGY1, AMMONIUM OVERLY SENSITIVE 1, AMOS1, ENHANCER OF VARIATION3, ETHYLENE-DEPENDENT GRAVITROPISM-DEFICIENT AND YELLOW-GREEN 1, EVR3
Immunogen:	KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from <i>Arabidopsis thaliana</i> EGY1 (AT5G35220).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum Peptide affinity form antibody available upon request at info@phytoab.com .
Reconstitution:	Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Research Use Only

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	59 kDa
Predicted Reactivity:	For more species homologues information, please contact tech support at tech@phytoab.com .

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.