

## Anti-Arogenate dehydratase 5, chloroplastic, N-terminal antibody

Catalog: PHY7542S

## **Product Information**

Description:	Rabbit polyclonal antibody
Background:	ADT5 is a plastid-localized arogenate dehydratase involved in phenylalanine
	biosynthesis. Not less than six genes encoding ADT were identified in the
	Arabidopsis genome: ADT1 [At1g11790]; ADT2 [At3g07630]; ADT3
	[At2g27820]; ADT4 [At3g44720]; ADT5 [At5g22630]; and ADT6 [At1g08250].
Synonyms:	ADT5, AROGENATE DEHYDRATASE 5
Immunogen:	KLH-conjugated synthetic peptide (14 aa from N terminal section) derived from
	Arabidopsis thaliana ADT5 (AT5G22630).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .
<b>Reconstitution:</b>	Reconstitution with 150 µl of sterile water.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ C$ as supplied.
	6 months, -20 to -70 $^\circ C$ under sterile conditions after reconstitution.
	1 month, 2 to 8 $^\circ\!\!\!\!\!^\circ \mathbb C$ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\!\mathbb{C}$ . Upon receipt, store it immediately at the
	temperature recommended above.

## **Application Information**


<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	46 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used

PhytoAB Inc.



for immunization is 100% homologues with the sequence in Brassica rapa, Brassica napus and 80-99% homologues with the sequence in Populus trichocarpa, Gossypium raimondii. For more species homologues information, please contact tech support at tech@phytoab.com.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 44 kDa. Electrophoresis: 12% SDS-PAGE Transfer: blotting to NC (nitrocellulose) membrane for 1 h. Blocking: 5% skim milk at RT or 4°C for 1 h. **Primary antibody:** 1:1000 dilution overnight at 4 °C. Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000). Detection: using chemiluminescence substrate and image were captured with CCD camera.

PHY7542S

Research Use Only