

Anti-Dihydrodipicolinate reductase-like protein CRR1, chloroplastic, C-terminal antibody

Catalog: PHY2822S

Product Information

| Description: | Rabbit polyclonal antibody |
|------------------------|--------------------------------------------------------------------------------------------------|
| Background: | CRR1 is essential for chloroplast NAD(P)H dehydrogenase activity, which is |
| | involved in electron transfer between PSII and PSI. |
| Synonyms: | CRR1, CHLORORESPIRATION REDUCTION 1 |
| Immunogen: | KLH-conjugated synthetic peptide (18 aa from C terminal section) derived from |
| | Arabidopsis thaliana CRR1 (AT5G52100). |
| Form: | Lyophilized |
| Quantity: | 150 µg |
| Purification: | Serum |
| | Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> . |
| Reconstitution: | Reconstitution with 150 µl of sterile water. |
| | "Note: please spin tube briefly prior to op <mark>ening it to avoid any losses that might</mark> |
| | occur from lyophilized material adhering to the cap or sides of the tube". |
| Stability & | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. |
| Storage: | 12 months from date of receipt, -20 to -70 $^\circ \! \mathbb C$ as supplied. |
| | 6 months, -20 to -70 $^\circ\!\!\mathbb{C}$ under sterile conditions after reconstitution. |
| | 1 month, 2 to 8 $^\circ$ under sterile conditions after reconstitution. |
| Shipping: | The product is shipped at 4 $^\circ\!\!\mathbb{C}$. Upon receipt, store it immediately at the |
| | temperature recommended above. |
| | |


Application Information

| Recommended Dilution: | Western Blot (1:1000-1:2000) |
|------------------------------|--------------------------------------------------------------------|
| | Note: Optimal dilutions/concentrations should be determined by the |
| | end user. |
| Expected / apparent MW: | 32 kDa |
| Predicted Reactivity: | Among species analyzed, the sequence of the synthetic peptide |
| | used for immunization is 100% homologues with the sequence in |
| | Brassica napus, Brassica rapa and 80-99% homologues with the |



sequence in Medicago truncatula, Glycine max, Cucumis sativus, Gossypium raimondii, Populus trichocarpa, Vitis vinifera. For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

PHY2822S

Research Use Only