

Anti-Auxin response factor 5, N-terminal antibody

Catalog: PHY3116A

Product Information

Description: Rabbit polyclonal antibody

Background: ARF5 is a transcription factor (IAA24) mediating embryo axis formation and

vascular development. It is similar to AUXIN RESPONSIVE FACTOR 1 (ARF1)

(AT1G59750) shown to bind to auxin responsive elements (AREs).

Synonyms: ARF5, IAA24, AUXIN RESPONSE FACTOR 5, INDOLE-3-ACETIC ACID

INDUCIBLE 24, MONOPTEROS, MP

Immunogen: KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from

Arabidopsis thaliana ARF5 (AT1G19850).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of sterile 1XPBS (PH=7.4).

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8° C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 100 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in *Brassica* napus, *Brassica* rapa, and 80-99% homologues with the sequence


in Populus trichocarpa, Gossypium raimondii.

Research Use Only



For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.