

## Anti-LRR receptor-like serine/threonine-protein kinase ERECTA antibody

Catalog: PHY1989A

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** ERECTA is a receptor protein kinase which is involved in specification of

organs originating from the shoot apical meristem. It contains a cytoplasmic protein kinase catalytic domain, a transmembrane region, and an extracellular leucine-rich repeat. ER has been identified as a quantitative trait locus for transpiration efficiency by influencing epidermal and mesophyll development, stomatal density and porosity of leaves. It has been implicated in resistance to

the bacterium Ralstonia solanacearum and to the necrotrophic fungus

Plectosphaerella cucumerina.

**Synonyms:** ERECTA, QRP1, QUANTITATIVE RESISTANCE TO PLECTOSPHAERELLA 1,

ER

**Immunogen:** KLH-conjugated synthetic peptide (18 aa from Central section) derived from

Arabidopsis thaliana ERECTA (AT2G26330).

Form: Lyophilized

**Quantity:** 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 μl of sterile 1XPBS (PH=7.4).

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.



## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected/apparent MW:** 107 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used for


immunization is 80-99% homologues with the sequence in Brassica

napus, Brassica rapa.

For more species homologues information, please contact tech

support at tech@phytoab.com.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 58 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.