

## Anti-NADH dehydrogenase subunit AT3G03070, mitochondrial antibody

Catalog: PHY3136A

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** Complex I is the largest protein complex of the oxidative phosphorylation

system in mitochondrial and it catalyzes NADH-quinone oxidoreduction.

Complex I represents the main entrance site for electrons into the respiratory electron transfer chain. In Arabidopsis, Complex I have at least 49 subunits and

AT3G03070 may be one of the subunit.

Synonyms: AT3G03070

Immunogen: KLH-conjugated synthetic peptide (16 aa from Central section) derived from

Arabidopsis thaliana NADH (AT3G03070).

Form: Lyophilized

**Quantity:** 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 µl of sterile 1XPBS (PH=7.4).

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 12 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in Brassica



napus, Brassica rapa, and 80-99% homologues with the sequence in Gossypium raimondii, Zea mays, Oryza sativa, Triticum aestivum, Hordeum vulgare, Panicum virgatum, Setaria viridis, Populus trichocarpa, Glycine max, Medicago truncatula, Cucumis sativus, Vitis vinifera, Spinacia oleracea, Nicotiana tabacum, Solanum lycopersicum, Solanum tuberosum.

For more species homologues information, please contact tech support at <a href="tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 40 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.