

Anti-Protection of telomeres protein 1b, N-terminal antibody

Catalog: PHY1869S

Product Information

Description:	Rabbit polyclonal antibody
Background:	Pot1 (protection of telomeres 1) is a single-stranded telomere binding protein
	that is essential for chromosome end protection and telomere length
	homeostasis. Arabidopsis has two Pot1-like proteins, dubbed POT1A
	(AT2G05210) and POT1B (AT5G06310).
Synonyms:	POT1B, ATPOT1B, PROTECTION OF TELOMERES 1B
Immunogen:	KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from
	Arabidopsis thaliana POT1B (AT5G06310).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .
Reconstitution:	Reconstitution with 150 µl of sterile water.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^{\circ}$ C as supplied.
	6 months, -20 to -70 $^\circ$ C under sterile conditions after reconstitution.
	1 month, 2 to 8 $^\circ\!\mathrm{C}$ under sterile conditions after reconstitution.
Shipping:	The product is shipped at $4^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the
	temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	53 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used


PhytoAB Inc.



for immunization is 80-99% homologues with the sequence in *Brassica napu*, *Brassica rapa*.

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 47 kDa. **Electrophoresis:** 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4 °C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

PHY1869S

Research Use Only