

Anti-Transcription factor MYB75, C-terminal antibody

Catalog: PHY1193A

Product Information

Description:	Rabbit polyclonal antibody
Background:	MYB75 is a putative MYB domain containing transcription factor involved in anthocyanin metabolism and radical scavenging. It is essential for the sucrose-mediated expression of the dihydroflavonol reductase gene. It interacts with JAZ proteins to regulate anthocyanin accumulation.
Synonyms:	MYB75, PAP1, ARABIDOPSIS THALIANA PRODUCTION OF ANTHOCYANIN PIGMENT 1, ATMYB75, ATPAP1, MYB DOMAIN PROTEIN 75, MYELOBLASTOSIS PROTEIN 75, PRODUCTION OF ANTHOCYANIN PIGMENT 1, SIAA1, SUC-INDUCED ANTHOCYANIN ACCUMULATION 1
Immunogen:	KLH-conjugated synthetic peptide (14 aa from C terminal section) derived from <i>Arabidopsis thaliana</i> MYB75 (AT1G56650).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of sterile 1XPBS (PH=7.4). "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	28 kDa

Research Use Only

Predicted Reactivity:

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.