

Anti-Glucocorticoid receptor antibody

Catalog: PHY6563A

Product Information

Description:	Rabbit polyclonal antibody
Background:	NR3C1
Synonyms:	NR3C1, Nr3c1
Immunogen:	KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from
	Rattus norvegicus NR3C1 (E9PT44).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of sterile 1XPBS (PH=7.4).
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	Note, please spin tube bleny phot to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	
Stability & Storage:	occur from lyophilized material adhering to the cap or sides of the tube".
	occur from lyophilized material adhering to the cap or sides of the tube". Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
	occur from lyophilized material adhering to the cap or sides of the tube". Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied.
	occur from lyophilized material adhering to the cap or sides of the tube". Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution.


Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	88 kDa
Predicted Reactivity:	For more species homologues information, please contact tech
	support at tech@phytoab.com.

Research Use Only



Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only