

Anti-Nitrate reductase [NADH] 1, N-terminal antibody

Catalog: PHY0047S

Product Information

Description: Rabbit polyclonal antibody

Background: Assimilatory nitrate reductase is an enzyme of the assimilative metabolism

involved in reduction of nitrate to nitrite. The nitrite is immediately reduced to

ammonia (probably via hydroxylamine) by the activity of nitrite reductase.

Plants contain 2 forms of NR: NADH-NR (most common form in plants and

algae, predominantly found in green tissues) and NAD(P)H-NR (uses NADH or

NADPH as the electron donor, constitutively expressed in plants at a low level).

NADH-NR is a homodimer of two identical subunits (100-115 kDa each, hold

together by a Mo-cofactor) each of them coded by up to three genes (NR1-3,

NIA1-NIA3).

Synonyms: NR1, GNR1, NITRATE REDUCTASE 1, NIA1

Immunogen: KLH-conjugated synthetic peptide (16 aa from N terminal section) derived from

Arabidopsis thaliana NR1 (AT1G77760).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the



end user.

Expected / apparent MW: 103 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 80-99% homologues with the sequence in

Brassica napus, Brassica rapa.

For more species homologues information, please contact tech

support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 49 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG

H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.