

Anti-Nuclear cap-binding protein subunit 2 antibody

Catalog: PHY2015A

Product Information

Description:	Rabbit polyclonal antibody
Background:	CBP20 is a nuclear cap-binding protein that forms a heterodimeric complex
	with ABH1 (AT2G13540) and is likely to participate in RNA metabolism. It
	recognizes and binds capped RNAs (m7GpppG-capped RNA) but requires
	ABH1/CBP80 to stabilize the movement of its N-terminal loop and lock the
	CBC into a high affinity cap-binding state with the cap structure.
Synonyms:	CBP20, ATCBP20, CAP-BINDING PROTEIN 20
Immunogen:	KLH-conjugated synthetic peptide (16 aa from Central section) derived from
	Arabidopsis thaliana CBP20 (AT5G44200).
Form:	Lyophilized
Quantity:	150 μg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of sterile 1XPBS (PH=7.4).
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70°C as supplied.
	6 months, -20 to -70°C under sterile conditions after reconstitution.
	1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the
	temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	30 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide
	used for immunization is 100% homologues with the sequence in


PhytoAB Inc.



Brassica rapa, Brassica napus, 80-99% homologues with the sequence in Gossypium raimondii, Panicum virgatum, Zea mays, Medicago truncatula, Glycine max, Populus trichocarpa, Solanum tuberosum, Solanum lycopersicum, Nicotiana tabacum, Vitis vinifera, Spinacia oleracea, Cucumis sativus, Oryza sativa, Setaria viridis, Sorghum bicolor, Triticum aestivum, Hordeum vulgareChlamydomonas reinhardtii. For more species homologues information, please contact tech

support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 47 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only