

Anti-Photosystem I assembly protein Ycf3 antibody

Catalog: PHY5444S

Product Information

Description:	Rabbit polyclonal antibody	
Background:	YCF3	
Synonyms:	YCF3	
Immunogen:	KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from	
	Synechocystis sp. PCC 6803 YCF3 (slr0823).	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Serum	
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .	
Reconstitution:	Reconstitution with 150 µl of sterile water.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70°C as supplied.	
	6 months, -20 to -70°C under sterile conditions after reconstitution.	
	1 month, 2 to 8°C under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the	
	temperature recommended above.	

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	20 kDa
Predicted Reactivity:	For more species homologues information, please contact tech
	support at tech@phytoab.com.

Research Use Only



Application Example

Example 1



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa. **Electrophoresis:** 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.


Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Example 2



PCC 6803-Thy: 12.5 μg, 25 μg and 50 μg thylakoid membrane protein from WT of *Synechocystis sp.* PCC 6803, respectively. **Electrophoresis:** 15% SDS-PAGE **Transfer:** blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

PHY5444S

Research Use Only