

Anti-Photosystem I assembly protein Ycf3 antibody

Catalog: PHY5447S

Product Information

| Description: | Rabbit polyclonal antibody |
|-----------------|--|
| Background: | YCF3 |
| Synonyms: | YCF3 |
| Immunogen: | Recombinant protein of YCF3 (1-173 aa) derived from Synechocystis sp. PCC |
| | 6803 slr0823. |
| Form: | Lyophilized |
| Quantity: | 150 μg |
| Purification: | Serum |
| Reconstitution: | Reconstitution with 150 µl of sterile water. |
| | · · · · · · · · · · · · · · · · · · · |
| | "Note: please spin tube briefly prior to opening it to avoid any losses that might |
| | |
| Stability & | "Note: please spin tube briefly prior to opening it to avoid any losses that might |
| | "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube". |
| Stability & | "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube". Use a manual defrost freezer and avoid repeated freeze-thaw cycles. |
| Stability & | "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube". Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. |
| Stability & | "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube". Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. |


Application Information

| Recommended Dilution: | Western Blot (1:1000-1:2000) |
|------------------------------|--|
| | Note: Optimal dilutions/concentrations should be determined by the |
| | end user. |
| Expected / apparent MW: | 20 kDa |
| Predicted Reactivity: | For more species homologues information, please contact tech |
| | support at <u>tech@phytoab.com</u> . |

Research Use Only



Application Example



PCC 6803-Thy: 12.5 µg, 25 µg and 50 µg thylakoid membrane protein from WT of *Synechocystis sp.* PCC 6803, respectively.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

PHY5447S



Research Use Only

PhytoAB Inc.