

Anti-Heat stress transcription factor A-1b antibody

Catalog: PHY7588S

Product Information

Description:	Rabbit polyclonal antibody	
-		
Background:	HSF3 is a putative transcription factor whose expression is not induced by heat	
	but whose stable overexpression leads to expression of HSP.	
Synonyms:	HSF3, ARABIDOPSIS HEAT SHOCK FACTOR 3, ARABIDOPSIS THALIANA	
	CLASS A HEAT SHOCK FACTOR 1B, ATHSF3, ATHSFA1B, CLASS A HEAT	
	SHOCK FACTOR 1B, HEAT SHOCK FACTOR 3, HSFA1B	
Immunogen:	KLH-conjugated synthetic peptide (15 aa from Central section) derived from	
	Arabidopsis thaliana HSF3 (AT5G16820).	
Form:	Lyophilized	
Quantity:	150 μg	
Purification:	Serum	
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .	
Reconstitution:	Reconstitution with 150 µl of sterile water.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70°C as supplied.	
	6 months, -20 to -70°C under sterile conditions after reconstitution.	
	1 month, 2 to 8°C under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the	
	temperature recommended above.	


Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	54 kDa
Predicted Reactivity:	For more species homologues information, please contact tech
	support at <u>tech@phytoab.com</u> .

Research Use Only



Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa. **Electrophoresis:** 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.



Research Use Only