

## Anti-C4-dicarboxylate transporter/malic acid transporter, N-terminal antibody

## **Product Information**

Description:	Rabbit polyclonal antibody
Background:	MTR_6g045200
Synonyms:	MTR_6g045200
Immunogen:	KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from
	<i>Medicago truncatula</i> MTR_6g045200.
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
<b>Reconstitution:</b>	Reconstitution with 150 µl of sterile 1XPBS (PH=7.4).
	"Note: please spin tube briefly prior to op <mark>ening it to avoid any losses that might</mark>
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ\!{ m C}$ as supplied.
	6 months, -20 to -70 $^\circ\!\!\mathbb{C}$ under sterile conditions after reconstitution.
	1 month, 2 to 8 $^{\circ}$ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\!\mathbb{C}$ . Upon receipt, store it immediately at the
	temperature recommended above.


## **Application Information**

<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	65 kDa
Predicted Reactivity:	For more species homologues information, please contact tech
	support at <u>tech@phytoab.com</u> .

Research Use Only



## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa. **Electrophoresis:** 12% SDS-PAGE **Transfer:** blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4°C for 1 h. **Primary antibody:** 1:1000 dilution overnight at 4°C. **Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.



**Research Use Only**