

Anti-C4-dicarboxylate transporter/malic acid transporter, C-terminal antibody

Catalog: PHY6453A

Product Information

Description:	Rabbit polyclonal antibody
Background:	MTR_6g045200
Synonyms:	MTR_6g045200
Immunogen:	KLH-conjugated synthetic peptide (16 aa from C terminal) derived from
	<i>Medicago truncatula</i> MTR_6g045200.
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of sterile 1XPBS (PH=7.4).
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ \!\! \mathbb{C}$ as supplied.
	6 months, -20 to -70 $^\circ\!\!\mathbb{C}$ under sterile conditions after reconstitution.
	1 month, 2 to 8 $^{\circ}$ C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\!{ m C}$. Upon receipt, store it immediately at the
	temperature recommended above.


Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	65 kDa
Predicted Reactivity:	For more species homologues information, please contact tech
	support at tech@phytoab.com.

Research Use Only



Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa. Electrophoresis: 12% SDS-PAGE Transfer: blotting to NC (nitrocellulose) membrane for 1 h. Blocking: 5% skim milk at RT or 4°C for 1 h. Primary antibody: 1:1000 dilution overnight at 4°C. Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000). Detection: using chemiluminescence substrate and image were

captured with CCD camera.

