

Anti-NADH dehydrogenase 6 antibody

Catalog: PHY2844A

Product Information

Description:	Rabbit polyclonal antibody
Background:	Complex I is the largest protein complex of the oxidative phosphorylation
	system in mitochondrial and it catalyzes NADH-quinone oxidoreduction.
	Complex I represents the main entrance site for electrons into the respiratory
	electron transfer chain. In Arabidopsis, Complex I have at least 49 subunits and
	NAD6 (ATMG00270) is one of the subunit.
Synonyms:	NAD6, NADH DEHYDROGENASE 6
Immunogen:	KLH-conjugated synthetic peptide of (18 aa from C terminal section) derived
	from Arabidopsis thaliana NAD6 (ATMG00270).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 μl of sterile 1XPBS (PH=7.4).
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70°C as supplied.
	6 months, -20 to -70°C under sterile conditions after reconstitution.
	1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the
	temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	24 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 100% homologues with the sequence in Solanum


Research Use Only



tuberosum, Triticum aestivum, Hordeum vulgare, Setaria viridis, Sorghum bicolor, Glycine max, Panicum virgatum, Nicotiana tabacum, Zea mays, Solanum lycopersicum, Cucumis sativus, Populus trichocarpa, Oryza sativa, Medicago truncatula, Spinacia oleracea, Brassica rapa, and 80-99% homologues with the sequence in Vitis vinifera, Gossypium raimondii.

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa. **Electrophoresis:** 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only