

Anti-Meiosis-specific protein ASY1 antibody

Catalog: PHY1170A

Product Information

Description:	Rabbit polyclonal antibody	
Background:	ASY1 protein is initially distributed as numerous foci throughout the chromatin.	
	It is required for normal meiosis in male and female gametophytes. And it plays	
	a crucial role in coordinating the activity of DMC1, a key member of the	
	homologous recombination machinery.	
Synonyms:	ASY1, ASYNAPTIC 1, ATASY1	
Immunogen:	KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from	
	Arabidopsis thaliana ASY1 (AT1G67370).	
Form:	Lyophilized	
Quantity:	150 μg	
Purification:	Immunogen affinity purified	
Reconstitution:	Reconstitution with 150 µl of sterile 1XPBS (PH=7.4).	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70°C as supplied.	
	6 months, -20 to -70°C under sterile conditions after reconstitution.	
	1 month, 2 to 8°C under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the	
	temperature recommended above.	

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	67 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 100% homologues with the sequence in <i>Cucumis</i>
	sativus, and 80-99% homologues with the sequence in Solanum
	tuberosum, Spinacia oleracea, Nicotiana tabacum, Gossypium


PhytoAB Inc.



raimondii, Solanum lycopersicum, Glycine max, Brassica napus, Brassica rapa, Medicago truncatula, Vitis vinifera, Populus trichocarpa, Triticum aestivum, Hordeum vulgare, Oryza sativa, Panicum virgatum, Setaria viridis, Sorghum bicolor, Zea mays, Sorghum bicolor.

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 40 kDa.
Electrophoresis: 12% SDS-PAGE
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).
Detection: using chemiluminescence substrate and image were captured

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only